


# Physics 221, April 13

## Key Concepts:

- Traveling sound waves
- Standing sound waves
- The Doppler effect

# Traveling sound waves



For a traveling sound wave in we have for the **pressure variations**

$$\Delta P(x,t) = \Delta P_{\max} \sin(kx - \omega t + \phi).$$

The corresponding **displacements from the equilibrium position** are

$$\Delta s(x,t) = \Delta s_{\max} \cos(kx - \omega t + \phi).$$

The **speed of sound in air** at standard temperature and pressure is **343** m/s, independent of the frequency or wavelength.

The **energy** carried by a sound wave is proportional to the square of its amplitude.


**Sound levels**  $\beta$  are measured using a logarithmic scale.

$$\beta = 10 \log_{10}(I/I_0)$$


The units of  $\beta$  are decibels (dB).  $I_0 = 10^{-12} \text{ W/m}^2$  is the reference intensity.

**MULTIPLY** the intensity by 10  $\Leftrightarrow$  **ADD** another 10 dB to the sound level

Which sounds travel the fastest through air?



1. higher pitch sound
2. lower pitch sound
3. louder sound
4. quieter sound
5. All sound travels at the same speed through air.



A pure sound notes from a sources make the molecules of air at a location vibrate simple harmonically in accordance with the equation


$$y_1 = 0.008 \sin (302 \pi t).$$

What is the frequency of the sound wave?

What is  $\omega$ ?

What is  $f$ ?

1. 604 Hz
2. 302 Hz
3. 96 Hz
4. 151 Hz



If the intensity of a 40 dB sound is increased to 80 dB, the intensity in  $\text{W}/\text{m}^2$  increases by a factor of

1. 2.
2. 4.
3. 40.
4.  $10^4$ .
5.  $10^{40}$ .

Hint:

MULTIPLY the intensity by 10  $\leftarrow = \rightarrow$


ADD another 10 dB to the sound level

Sound level examples:

10 dB: Rustling or falling leaves

50 dB: Conversation.


100 dB: Hearing damage after about 15 minutes.



# Beats

Intensity modulations are produced when waves of slightly different frequencies are superimposed.

The beat frequency is equal to the difference frequency  $|f_1 - f_2|$ .



## Demonstration:


A speaker box containing two oscillators, one fixed and the other variable.

By tuning the variable oscillator we can produce audio beats.

Two flutists are tuning up. If the conductor hears the beat frequency increasing, are the two flute frequencies getting closer together or farther apart?




1. Closer
2. Farther apart
3. No way to tell



# Interference

**Demonstration:** Two speakers and a sine wave generator. The speakers are driven by the same signal and can move in or 180 degrees out of phase.

Like all waves, two or more sound waves traveling through the same medium will interfere. We can have constructive and destructive interference.



Assume  $s_1 = 5$  m and  $s_2 = 4$  m. The path difference  $\Delta = 1$  m. If  $\Delta = m \cdot \lambda$  with  $m = \text{integer}$ , then we have constructive interference.


**Constructive interference:**  $\lambda = \Delta/m$ ,  $f = v/\lambda = mv/\Delta$ .

**Destructive interference:**  $\lambda = \Delta/(m + \frac{1}{2})$ ,  $f = v/\lambda = (m + \frac{1}{2})v/\Delta$ .




In the figure below, two point sources  $S_1$  and  $S_2$ , which are in phase, emit identical sound waves of wavelength 2.0 m.

If  $L_1 = 39 \text{ m}$  and  $L_2 = 36 \text{ m}$ , what type of interference occurs at point P?



1. Constructive interference
2. Destructive interference
3. Beats
4. No interference at all



Constructive interference:  $\lambda = \Delta/m$ .

Destructive interference:  $\lambda = \Delta/(m + \frac{1}{2})$ .

$m$  is an integer.


Here  $\lambda = 2 m = \Delta/1.5 = \Delta/(1 + \frac{1}{2})$ .

# Standing sound waves

We can create a standing wave in a tube, which is open on both ends, and in a tube, which is open on one end and closed on the other end. **Open and closed ends** reflect waves differently. Displacement variations are plotted.


Fundamental

$$\lambda = 2L \quad f = v / 2L$$




Fundamental

$$\lambda = 4L \quad f = v / 4L$$




Second harmonic

$$\lambda = L \quad f = v / L$$



Third harmonic

$$\lambda = 4/3 L \quad f = 3v / 4L$$




**Demonstration:** Singing Tubes


Organ pipe B is open at both ends and is **half as long** as organ pipe A, which is open at one end, as shown.

What is

- (i) the ratio of their fundamental wavelengths  $\lambda_A : \lambda_B$  and
- (ii) the ratio of their fundamental frequencies  $f_A : f_B$  ?



- 1. (i) 1:1 (ii) 1:1
- 2. (i) 1:2 (ii) 1:2
- 3. (i) 2:1 (ii) 2:1
- 4. (i) 1:4 (ii) 4:1
- 5. (i) 4:1 (ii) 1:4



# Doppler effect

The perceived pitch of a sound wave changes if the observer or the source is moving.

If both source and observer are in motion, then the apparent frequency of the sound wave reaching the observer is

$$f = f_0(v - v_{\text{obs}})/(v - v_s)$$

$v$  = velocity of sound

$v_o$  = velocity component of the observer in the direction of  $v$

$v_s$  = velocity component of the source in the direction of  $v$




**Demonstration:** The Doppler Rocket


As a sound source moving with constant velocity approaches and then moves past a stationary observer, the observer will hear



1. a steady rise in pitch.
2. a sudden drop in pitch.
3. a rise in pitch, then a drop in pitch.
4. a drop in pitch, then a rise in pitch.



When a car is at rest, its horn emits a sound wave of wavelength 0.55 m. A person standing in the middle of the street hears the horn with a frequency of 560 Hz. If the speed of sound is 330 m/s, should the person jump out of the way?



1. Yes

2. **No**

